

Edge-transitive embeddings of complete graphs

Gareth A. Jones* 

School of Mathematics, University of Southampton, Southampton SO17 1BJ, UK

Received 9 August 2019, accepted 14 April 2020, published online 13 August 2021

Abstract

Building on earlier work of Biggs, James, Wilson and the author and on the Graver–Watkins description of the 14 classes of edge-transitive maps, we complete the classification of the edge-transitive embeddings of complete graphs by classifying their non-regular embeddings in non-orientable surfaces and their embeddings with non-empty boundary.

DEDICATED TO THE MEMORY OF BRANKO GRÜNBAUM

Keywords: Edge-transitive map, complete graph, Biggs map, James map.

Math. Subj. Class.: 05C10, 20B25.

1 Introduction

The purpose of this paper is to complete the classification of edge-transitive embeddings of complete graphs (see Theorem 1.8) by classifying the non-regular embeddings in non-orientable surfaces in Theorem 1.7 and the embeddings with non-empty boundary in Theorem 6.1. First we review the earlier results in this area.

In 1987 Grünbaum and Shephard [6] published an influential paper on edge-transitive planar graphs, extending earlier work on finite edge-transitive graphs to the infinite case. An important consequence of this was the monograph [4] by Graver and Watkins; this included a partition of all edge-transitive maps \mathcal{M} into 14 classes, which can be interpreted as corresponding to the 14 possibilities for $\mathcal{M}/\text{Aut } \mathcal{M}$. These classes were studied further by Širáň, Tucker and Watkins [20], by Orbanič, Pellicer, Pisanski and Tucker [17] and by the author [12], investigating the possible surfaces and automorphism groups for maps in

*The author is grateful to the referee for some suggestions which have significantly improved the presentation of this paper.

E-mail address: G.A.Jones@maths.soton.ac.uk (Gareth A. Jones)

the various classes, mainly in the finite case. The aim of this note is to approach edge-transitive maps from the point of view of their embedded graphs, specifically when these are complete graphs.

A map on a surface is *regular* (sometimes called *fully regular*) or *edge-transitive* if its automorphism group acts transitively on vertex-edge-face flags or edges respectively. A map on an orientable surface is *orientably regular* if its orientation-preserving automorphism group acts transitively on arcs, and *chiral* if it has no orientation-reversing automorphisms, so that it is not isomorphic to its mirror image. Thus an orientably regular map is either regular or chiral, but not both. The regular and orientably regular embeddings of complete graphs are all known, as are their edge-transitive orientable embeddings. This paper will deal with the remaining cases, the non-regular embeddings in non-orientable surfaces, together with those in surfaces with non-empty boundary. (Unless otherwise stated, as in Section 6, all maps considered from now on are assumed to be without boundary.)

In 1971 Biggs [1], building on earlier work of Heffter [7], proved:

Theorem 1.1. *The complete graph K_n has an orientably regular embedding if and only if n is a prime power.*

The maps Biggs constructed to prove that this condition is sufficient are Cayley maps (see [5, §1.2.4]) $\mathcal{M}_n(c)$ for the additive groups of finite fields \mathbb{F}_n ; in each case the generating set is the multiplicative group $\mathbb{F}_n^* = \mathbb{F}_n \setminus \{0\}$, taken in the cyclic order $1, c, c^2, \dots, c^{n-2}$ where c is a primitive element of \mathbb{F}_n , that is, a generator for the cyclic group \mathbb{F}_n^* . They are chiral for all $n \geq 5$.




Figure 1: The Biggs maps $\mathcal{M}_5(2)$ and $\mathcal{M}_7(3)$

Example 1.2. For $n = 4$ we have $\mathbb{F}_4 = \{0, 1, c, c^2 = c^{-1} = c + 1\}$, and the corresponding map $\mathcal{M}_4(c)$ is the tetrahedral map on the sphere. The Biggs maps $\mathcal{M}_5(2)$ and $\mathcal{M}_7(3)$ are shown in Figure 1, with opposite sides of the outer square and hexagon identified to form the torus maps $\{4, 4\}_{1,2}$ and $\{3, 6\}_{1,2}$ in the notation of [3, §8.3 and §8.4]. Their mirror images are the maps $\mathcal{M}_5(3) \cong \{4, 4\}_{2,1}$ and $\mathcal{M}_7(5) \cong \{3, 6\}_{2,1}$.

In 1985 James and the author [11] proved that the Biggs maps $\mathcal{M}_n(c)$ are the only orientably regular embeddings of complete graphs:

Theorem 1.3. *A map \mathcal{M} is an orientably regular embedding of K_n if and only if $\mathcal{M} \cong \mathcal{M}_n(c)$ for some primitive element c of \mathbb{F}_n . Moreover, $\mathcal{M}_n(c)$ and $\mathcal{M}_n(c')$ are isomorphic (as oriented maps) if and only if c and c' are equivalent under a field automorphism of \mathbb{F}_n .*

It follows that $n = p^e$ for some prime p and there are, up to isomorphism, $\phi(n - 1)/e$ orientably regular embeddings of K_n , one for each orbit of the Galois group of \mathbb{F}_n (isomorphic to C_e , generated by the Frobenius automorphism $t \mapsto t^p$) on the $\phi(n - 1)$ primitive elements of the field. The orientation-preserving automorphism group of $\mathcal{M}_n(c)$ is the affine group $AGL_1(\mathbb{F}_n)$; this map is regular if and only if $n = 2, 3$ or 4 , in which case the full automorphism group is isomorphic to $V_4 \cong C_2 \times C_2$, $D_6 \cong S_3 \times C_2$ or S_4 .

If $n \geq 3$ the Petrie dual $P(\mathcal{M}_n(c))$ of $\mathcal{M}_n(c)$ is a non-orientable edge-transitive embedding of K_n , with the same automorphism group as $\mathcal{M}_n(c)$.

In [9] James classified the non-orientable regular embeddings of complete graphs (see also [22] for an independent proof due to Wilson):

Theorem 1.4. *The non-orientable regular embeddings of complete graphs K_n are the maps $\{6, 2\}_3$, $\{4, 3\}_3$, $\{3, 5\}_5$ and $\{5, 5\}_3$ of characteristic 1, 1, 1 and -3 for $n = 3, 4, 6$ and 6.*




Figure 2: Regular embeddings of K_n on the projective plane, $n = 3, 4, 6$

Here the notation $\{p, q\}_r$, from [3, §8.6 and Table 8], denotes the largest map of type $\{p, q\}$ with Petrie length r . The first three of these maps, on the real projective plane, are the antipodal quotients of a hexagon, a cube and an icosahedron on the sphere (see Figure 2, where antipodal boundary points of the discs are identified). The first two are the Petrie duals of the unique regular embeddings $\mathcal{M}_n(c)$ of K_3 and K_4 on the sphere, with the same automorphism groups D_6 and S_4 , while the last two are a Petrie dual pair with automorphism group $PSL_2(5) \cong A_5$.

In [10] James extended Theorem 1.3 to a classification of the orientable edge-transitive embeddings of K_n . If $3 < n = p^e \equiv 3 \pmod{4}$ where p is prime, and c is a primitive element of \mathbb{F}_n , let $\mathcal{M}_n(c, j)$ be the Cayley map for \mathbb{F}_n with generating set \mathbb{F}_n^* , where now the cyclic ordering is

$$1, c^j, c^2, c^{j+2}, c^4, c^{j+4}, \dots, c^{n-3}, c^{j+n-3}$$

for some odd element $j \in \mathbb{Z}_{n-1} \setminus \{1\}$. (Taking $j = 1$ gives the orientably regular Biggs map $\mathcal{M}_n(c)$.)

Theorem 1.5. *A map \mathcal{M} is an orientable edge-transitive embedding of K_n , which is not orientably regular, if and only if $\mathcal{M} \cong \mathcal{M}_n(c, j)$ for some n, c and j as above. As oriented maps, $\mathcal{M}_n(c, j)$ and $\mathcal{M}_n(c', j')$ are isomorphic if and only if c and c' are equivalent under $\text{Gal } \mathbb{F}_n$ and $j' \equiv j$ or $2 - j \pmod{n - 1}$.*

These James maps $\mathcal{M}_n(c, j)$ have automorphism group $AHL_1(\mathbb{F}_n)$, the unique subgroup of index 2 in $AGL_1(\mathbb{F}_n)$; their Petrie duals are non-orientable edge-transitive embeddings of K_n . Each map $\mathcal{M}_n(c, j)$ is chiral, with mirror image $\mathcal{M}_n(c^{-1}, 2 - j) \cong \mathcal{M}_n(c^{-1}, j)$.




Figure 3: An edge-transitive embedding $\mathcal{M}_7(5, 5)$ of K_7

Example 1.6. The James map $\mathcal{M}_7(5, 5)$ is shown in Figure 3, taken from [10]. Two of the identifications of sides of the outer 14-gon are indicated by the letters A and B ; the others can be found by C_7 rotational symmetry. The vertices are identified with the elements $0, 1, \dots, 6$ of \mathbb{F}_7 , in clockwise order. There are seven triangular faces and three heptagons. The mirror image $\mathcal{M}_7(3, 3)$ of this map is given by reflection in the horizontal axis. Each of these maps is a representation of the Fano plane $\mathbb{P}^2(\mathbb{F}_2)$, with the vertices as its points and the triples incident with its triangular faces forming its lines. (Similar representations of this and other finite projective planes are discussed by Singerman in [19].)

These two maps \mathcal{M} can be drawn as follows on Klein’s Riemann surface (or quartic curve) \mathcal{K} of genus 3 with automorphism group $PSL_2(7)$ (see [15], for instance), so that $\text{Aut } \mathcal{M} (\cong C_7 \times C_3)$ is contained in $\text{Aut } \mathcal{K}$. Each of the eight Sylow 7-subgroups $S \cong C_7$ of $\text{Aut } \mathcal{K}$ preserves three of the 24 faces of the $\text{Aut } \mathcal{K}$ -invariant tessellation $\mathcal{T} = \{7, 3\}_8$ of \mathcal{K} . These three faces are incident with 21 of the 56 vertices of \mathcal{T} , and there are 21 more vertices adjacent in \mathcal{T} to these. This leaves 14 vertices of \mathcal{T} , at graph-theoretic distance 2 from the S -invariant faces, forming two orbits under S . The vertices in each orbit, joined pairwise by geodesics, determine a chiral pair of maps $\mathcal{M} \cong \mathcal{M}_7(5, 5)$ and $\mathcal{M}_7(3, 3)$, each




Figure 4: Vertices of $\mathcal{M}_7(5, 5)$ and $\mathcal{M}_7(3, 3)$ on Klein's surface

invariant under the normaliser $\text{Aut } \mathcal{M} = S \rtimes C_3$ of S in $\text{Aut } \mathcal{K}$. Figure 4 shows part of \mathcal{T} , with the central face invariant under the group S generated by rotation through $2\pi/7$, and the orbits of S on vertices of \mathcal{T} yielding these two maps indicated in black and white. The vertices of each map are the centres of the triangular faces of the other. The three heptagonal faces of each map each contain one of the S -invariant faces of \mathcal{T} .

In order to complete the classification of edge-transitive embeddings of complete graphs, it remains for us to deal with the non-regular non-orientable cases. The main result of this paper, proved in Section 4 after some preparatory work in Sections 2 and 3, is as follows:

Theorem 1.7. *A map \mathcal{M} is a non-orientable non-regular edge-transitive embedding of a complete graph K_n if and only if \mathcal{M} is isomorphic to the Petrie dual of a Biggs map $\mathcal{M}_n(c)$ for $n \geq 5$ or of a James map $\mathcal{M}_n(c, j)$ for $n \geq 7$.*

As an immediate corollary we have the following classification of the edge-transitive embeddings of complete graphs, showing that they are simply the maps listed above:

Theorem 1.8. *A map \mathcal{M} is an edge-transitive embedding of a complete graph K_n if and only if \mathcal{M} is isomorphic to a Biggs map $\mathcal{M}_n(c)$ or its Petrie dual, a James map $\mathcal{M}_n(c, j)$ or its Petrie dual, or one of the Petrie dual pair $\{3, 5\}_5$ and $\{5, 5\}_3$ of non-orientable regular embeddings of K_6 .*

A detailed description of these maps is given in Section 5. Edge-transitive embeddings of K_n in surfaces with boundary are classified in Section 6.

Finally, we note that by results of Korzhik and Voss [14], vertex-transitive embeddings of K_n are much more abundant and harder to classify than its edge-transitive embeddings: for example, any cyclic ordering of the non-identity elements of a group of order n will yield such a Cayley map for that group. Nevertheless, Li gives a good description of the possibilities in [16].

2 Algebraic theory of maps

Here we sketch the algebraic theory of maps developed in more detail elsewhere: see [13], for example, and [5] for further background in topological graph theory.

Each map \mathcal{M} (possibly non-orientable or with non-empty boundary) determines a permutation representation of the group

$$\Gamma = \langle R_0, R_1, R_2 \mid R_i^2 = (R_0R_2)^2 = 1 \rangle \cong V_4 * C_2$$

on the set Φ of flags $\phi = (v, e, f)$ of \mathcal{M} , where v, e and f are a mutually incident vertex, edge and face. For each $\phi \in \Phi$ and each $i = 0, 1, 2$, there is at most one flag $\phi' \neq \phi$ with the same j -dimensional components as ϕ for each $j \neq i$ (possibly none if ϕ is a boundary flag). Define r_i to be the permutation of Φ transposing each ϕ with ϕ' if the latter exists, and fixing ϕ otherwise, in which case the incident edge is called a *free edge* or *semi-edge*. (See Figures 5 and 6 for these two cases. In Figure 6 the broken line represents part of the boundary of the map.) Since $r_i^2 = (r_0r_2)^2 = 1$ there is a permutation representation

$$\theta : \Gamma \rightarrow G := \langle r_0, r_1, r_2 \rangle \leq \text{Sym } \Phi$$

of Γ on Φ , given by $R_i \mapsto r_i$.




Figure 5: Generators r_i of G acting on a flag $\phi = (v, e, f)$.




Figure 6: Flags fixed by r_0, r_1 and r_2 .

Conversely, any permutation representation of Γ on a set Φ determines a map \mathcal{M} in which the vertices, edges and faces are identified with the orbits on Φ of the subgroups $\langle R_1, R_2 \rangle \cong D_\infty$, $\langle R_0, R_2 \rangle \cong V_4$ and $\langle R_0, R_1 \rangle \cong D_\infty$, incident when they have non-empty intersection.

The map \mathcal{M} is connected if and only if Γ acts transitively on Φ , as we will always assume. In this case the stabilisers in Γ of flags $\phi \in \Phi$ form a conjugacy class of subgroups $M \leq \Gamma$, called *map subgroups*.

The map \mathcal{M} is finite (has finitely many flags) if and only if M has finite index in Γ , and it has non-empty boundary if and only if M contains a conjugate of some R_i , or equivalently some r_i has a fixed point in Φ . In particular, \mathcal{M} is orientable and with empty boundary if and only if M is contained in the even subgroup Γ^+ of index 2 in Γ , consisting of the words of even length in the generators R_i .

The *automorphism group* $A = \text{Aut } \mathcal{M}$ of \mathcal{M} is the centraliser of G in $\text{Sym } \Phi$. Then $A \cong N/M$ where $N := N_\Gamma(M)$ is the normaliser of M in Γ . The map \mathcal{M} is called *regular* if A is transitive on Φ , or equivalently G is a regular permutation group, that is, M is normal in Γ ; in this case $A \cong G \cong \Gamma/M$, and one can identify Φ with G , so that A and G are the left and right regular representations of G . The map \mathcal{M} is *edge-transitive* if A acts transitively on its edges, or equivalently $\Gamma = NE$ where $E := \langle R_0, R_2 \rangle \cong V_4$.

The (classical) dual $D(\mathcal{M})$ of \mathcal{M} corresponds to the image of M under the automorphism δ of Γ fixing R_1 and transposing R_0 and R_2 . The Petrie dual $P(\mathcal{M})$ embeds the same graph as \mathcal{M} , but the faces are transposed with Petrie polygons, closed zig-zag paths turning alternately first right and first left at the vertices of \mathcal{M} ; this operation P corresponds to the automorphism π of Γ transposing R_0 with R_0R_2 and fixing R_1 and R_2 . Both of these operations D and P preserve automorphism groups and (full) regularity, but D may change the embedded graph, and P may change the underlying surface; in particular, if \mathcal{M} is orientable, then $P(\mathcal{M})$ is orientable if and only if the embedded graph is bipartite. The group $\Omega = \langle D, P \rangle$ of map operations generated by D and P , introduced by Wilson in [21], is isomorphic to S_3 , permuting vertices, faces and Petrie polygons; it corresponds to the outer automorphism group $\text{Out } \Gamma \cong \text{Aut } E \cong S_3$ of Γ acting on maps by permuting conjugacy classes of map subgroups [13].

3 Edge-transitive maps

In 1997 Graver and Watkins [4] partitioned the edge-transitive maps \mathcal{M} into 14 classes, distinguished by the isomorphism class of the quotient map $\mathcal{M}/\text{Aut } \mathcal{M}$; in that year, Wilson [23] gave a similar classification. These classes T correspond to the 14 isomorphism classes of maps $\mathcal{N}(T)$ with one edge, shown in Figure 7, or equivalently to the 14 conjugacy classes of *parent groups*, subgroups $N = N(T)$ of Γ satisfying $\Gamma = NE$ (see [12, §4]).

The maps in Figure 7 are all on the closed disc, apart from $\mathcal{N}(2^P \text{ex})$, $\mathcal{N}(5)$ and $\mathcal{N}(5^*)$ on the sphere, $\mathcal{N}(4^P)$ on the Möbius band and $\mathcal{N}(5^P)$ on the real projective plane. The 14 edge-transitive classes include class 1, consisting of the regular maps, and class 2^Pex , consisting of the chiral (non-regular) orientably regular maps. Each map \mathcal{M} in class T , with automorphism group A , is a regular covering by A of the basic map $\mathcal{N}(T)$ for that class; it corresponds to a map subgroup $M \leq \Gamma$ with $N_\Gamma(M) = N(T)$ and $N(T)/M \cong A$. The six rows of Figure 7 correspond to the orbits of Ω on the 14 classes: the duals of the maps in class 2 form class 2^* , while the Petrie duals of the latter form class 2^P ; similar remarks apply to the classes 2ex , 4 and 5, whereas the classes 1 and 3 are invariant under Ω . Thus, being chiral and vertex- but not face-transitive, $\mathcal{M}_n(c, j)$ is in class 5^* , so $P(\mathcal{M}_n(c, j))$ is in class 5^P .

The Reidemeister-Schreier process, applied to the inclusions $N(T) \leq \Gamma$, gives the




Figure 7: The basic maps $\mathcal{N}(T)$ for the 14 edge-transitive classes T

following presentations and free product decompositions:

$$\begin{aligned}
 N(1) &= \Gamma = \langle R_0, R_1, R_2 \mid R_i^2 = (R_0 R_2)^2 = 1 \rangle \cong V_4 * C_2, \\
 N(2) &= \langle S_1 = R_1, S_2 = R_1^{R_0}, S_3 = R_2 \mid S_1^2 = S_2^2 = S_3^2 = 1 \rangle \cong C_2 * C_2 * C_2, \\
 N(2 \text{ ex}) &= \langle S_1 = R_2, S = R_0 R_1 \mid S_1^2 = 1 \rangle \cong C_2 * C_\infty, \\
 N(3) &= \langle S_0 = R_1, S_1 = R_1^{R_0}, S_2 = R_1^{R_2}, S_3 = R_1^{R_0 R_2} \mid S_i^2 = 1 \rangle \cong C_2 * C_2 * C_2 * C_2, \\
 N(4) &= \langle S_1 = R_1, S_2 = R_1^{R_2}, S = (R_1 R_2)^{R_0} \mid S_1^2 = S_2^2 = 1 \rangle \cong C_2 * C_2 * C_\infty, \\
 N(5) &= \langle S = R_1 R_2, S' = S^{R_0} \mid - \rangle \cong C_\infty * C_\infty \cong F_2.
 \end{aligned}$$

(Here F_2 denotes a free group of rank 2.) Applying elements of $\text{Aut } E \cong \Omega$, permuting R_0, R_2 and $R_0 R_2$, gives presentations for the other (isomorphic) parent groups in each orbit of Ω .

4 Proof of Theorem 1.7

It is easy to see that $P(\mathcal{M}_n(c))$ and $P(\mathcal{M}_n(c, j))$ have the stated properties.

Conversely, if \mathcal{M} is a non-regular non-orientable edge-transitive embedding of K_n , then it is in one of the 10 edge-transitive classes $T \neq 1, 2^P \text{ex}, 5$ or 5^* , since the maps in class 1 are regular, while those in the other three classes are orientable. The cases $n \leq 3$ are easily dealt with, so assume that $n \geq 4$. If $T = 2^* \text{ex}$ or 5^P then $\mathcal{M} = P(\mathcal{M}')$ for some edge-transitive embedding \mathcal{M}' of K_n in class 2^Pex or 5^* ; as such, \mathcal{M}' is orientable, so by Theorems 1.3 and 1.5 it is isomorphic to a map $\mathcal{M}_n(c)$ or $\mathcal{M}_n(c, j)$ respectively as it is or is not orientably regular. This leaves only $T = 2, 2^*, 2^P, 2 \text{ex}, 3, 4, 4^*$ and 4^P , so it is sufficient to eliminate these cases.

The edges of K_n may be identified with the distinct pairs of vertices, so edge-transitivity implies that the group $A = \text{Aut } \mathcal{M}$ permutes these pairs transitively, that is, it acts 2-homogeneously on the vertices. It permutes the vertices faithfully since $n \geq 4$, and transitively since K_n is not bipartite; the latter property rules out classes $T = 2, 3$ and 4 , since the corresponding maps $\mathcal{N}(T)$ have two vertices, so only the classes $T = 2^*, 2^P, 2 \text{ex}, 4^*$ and 4^P remain. If A has odd order then, as a quotient of the group $N(T)$ with a free product decomposition given above, it must be cyclic; however, a cyclic group cannot be 2-homogeneous of degree $n \geq 4$, so A has even order. It thus has an element transposing two vertices, so it is 2-transitive on the vertices.

Each edge of \mathcal{M} is therefore reversed by some automorphism (a reflection or half turn), so the single edge of $\mathcal{N}(T)$ must be free, ruling out the classes $T = 2 \text{ex}, 4^*$ and 4^P . Only the classes 2^* and 2^P remain, with A sharply 2-transitive on the vertices (since \mathcal{M} is not regular), and the stabiliser of an edge generated by a reflection or a half-turn respectively, transposing its two incident vertices. Zassenhaus [24] showed that any sharply 2-transitive finite group can be identified with $AGL_1(F)$ acting naturally on a near-field F . Since \mathcal{M} is a map, the stabiliser A_0 of the vertex 0 must be a cyclic or dihedral group of order $n - 1$, acting regularly on the set $F \setminus \{0\}$ of neighbours of 0. Now A acts on the vertices as a Frobenius group $F \rtimes A_0$, so A_0 is a Frobenius complement; these contain at most one involution (see [8, Satz V.8.18(a)] or [18, Theorem 18.1(iii)]), so they cannot be dihedral, and hence A_0 is cyclic. Since $T = 2^*$ or 2^P , $N(T)$ is generated by involutions, and hence so are its epimorphic images A and A_0 , giving $n - 1 \leq 2$, a contradiction. \square

5 Summary of properties of the maps

The following basic properties of the maps classified in Theorem 1.8 are taken from [1, 9, 10, 11] or have been demonstrated earlier in this paper.

For each prime power $n = p^e$ there are $\phi(n - 1)/e$ Biggs maps $\mathcal{M}_n(c)$, all orientable. The mirror image of $\mathcal{M}_n(c)$ is $\mathcal{M}_n(c^{-1})$. For each $n = 2, 3, 4$ the unique map $\mathcal{M}_n(c)$ is regular (in class 1), of type $\{2, 1\}_2, \{3, 2\}_6$ or $\{3, 3\}_4$ and genus 0, with automorphism group V_4, D_6 or S_4 . For $n \geq 5$ these maps are orientably regular (in class 2^Pex) with automorphism group $AGL_1(\mathbb{F}_n)$; they have type $\{m, n - 1\}_{2p}$ where $m = (n - 1)/2$ or $n - 1$ as $n \equiv 3 \pmod{4}$ or not, and have genus $(n^2 - 7n + 4)/4$ or $(n - 1)(n - 4)/4$ respectively.

The maps $P(\mathcal{M}_n(c))$ for $n \geq 3$ are non-orientable, with the same automorphism group as $\mathcal{M}_n(c)$; they are in class 1 or 2^*ex as $n \leq 4$ or $n \geq 5$. They have type $\{2p, n - 1\}_m$

where m is as above, and characteristic

$$\chi = n \left(1 - \frac{n-1}{2} + \frac{n-1}{2p} \right).$$

If $3 < n = p^e \equiv 3 \pmod{4}$, there are $(n-3)\phi(n-1)/4e$ James maps $\mathcal{M}_n(c, j)$, all orientable, with mirror image $\mathcal{M}_n(c^{-1}, 2-j)$. They are in class 5^* , with automorphism group $AHL_1(\mathbb{F}_n)$ consisting of the transformations $v \mapsto av + b$ in $AGL_1(\mathbb{F}_n)$ such that a is a non-zero square in \mathbb{F}_n . This group has two orbits on the faces: if $j, 2-j \not\equiv (n-1)/2 \pmod{n-1}$ they have cardinality $n(n-1, j)$ and $n(n-1, 2-j)$ and the genus is $\frac{n}{4}((n-3) - 2(n-1, j) - 2(n-1, 2-j)) + 1$, whereas if j or $2-j \equiv (n-1)/2 \pmod{n-1}$ they have cardinality n and $n(n-1)/2p$ and the genus is $(n-1)(n(p-1) - 4p)/4p$. There is a single orbit of $n(n-1)/l$ Petrie polygons, each polygon having length $l = 2(n-1)/(n-1, 2(j-1))$.

The maps $P(\mathcal{M}_n(c, j))$ are non-orientable, with automorphism group $AHL_1(\mathbb{F}_n)$; they are in class 5^P , and have type $\{l, n-1\}$ and characteristic

$$\chi = n \left(1 - \frac{n-1}{2} + \frac{n-1}{l} \right).$$

For $n = 6$ the Petrie dual pair $\{3, 5\}_5$ and $\{5, 5\}_3$ are non-orientable, with $\chi = 1$ and -3 ; they are regular, with automorphism group $PSL_2(5) \cong A_5$.

These maps \mathcal{M} are all vertex-transitive. Indeed, in all cases $\text{Aut } \mathcal{M}$ has a subgroup acting regularly on the vertices, so that \mathcal{M} is a Cayley map: when n is a prime power there is a unique such subgroup, namely the additive group of \mathbb{F}_n acting by translations, whereas when $n = 6$ there is a conjugacy class of ten such subgroups, isomorphic to S_3 . Apart from the James maps $\mathcal{M}_n(c, j)$, these maps are all face-transitive. They are all arc-transitive with the exception of the James maps and their Petrie duals.

6 Edge-transitive embeddings with boundary

Here we will classify the edge-transitive maps \mathcal{M} which embed complete graphs K_n in surfaces with non-empty boundary $\partial\mathcal{M}$. (A general theory of maps with boundary was developed by Bryant and Singerman in [2].)




Figure 8: Edge-transitive embeddings of K_n with boundary, $n = 2, 3$

There are three maps each for $n = 2$ and 3, shown in Figure 8, and none for $n \geq 4$. The first five are on the closed disc, while the last is on the Möbius band. The first two embeddings of K_2 are in class 1, while the third is in class 2; their automorphism groups are isomorphic to C_2 , V_4 and C_2 . The three embeddings of K_3 , all with automorphism group S_3 , are in classes 1, 2^* and 2^P respectively. The last two form a Petrie dual pair.

Theorem 6.1. *The only edge-transitive maps which embed complete graphs K_n in surfaces with non-empty boundary are the six shown in Figure 8.*

Proof. Let \mathcal{M} be such a map. If $n = 2$ then \mathcal{M} has one edge, so it is one of the 14 basic maps $\mathcal{N}(T)$ with one edge shown in Figure 7. By inspection, $\mathcal{N}(T)$ embeds K_2 if and only if $T = 2, 3, 4$ or 5. Only the first three of these have non-empty boundary, giving the maps in the first row of Figure 8. We may therefore assume that $n \geq 3$. It follows that by edge-transitivity $\text{Aut } \mathcal{M}$ acts transitively (in fact, 2-homogeneously) on the vertices.

Since $\partial\mathcal{M} \neq \emptyset$, some flag ϕ of \mathcal{M} must be fixed by some r_i for $i = 0, 1, 2$. If $i = 0$ then \mathcal{M} has a free edge (see Figure 6), contradicting the fact that it embeds a complete graph. If $i = 2$ then the corresponding edge, and hence every edge, is contained in $\partial\mathcal{M}$; thus each vertex has valency at most 2, so $n = 3$ and we have the first graph in the second row of Figure 8.

We may therefore assume that only r_1 fixes a flag. By vertex-transitivity, every vertex is in $\partial\mathcal{M}$, with two of its $2(n - 1)$ incident flags fixed by r_1 . Thus the proportion of flags fixed by r_1 is $1/(n - 1)$. By edge-transitivity, each edge is incident with the same number of flags fixed by r_1 , which must be 1, 2, 3 or 4, so $1/(n - 1) = 1/4, 1/2, 3/4$ or 1 and hence $n = 5, 3, 7/3$ or 2. The last two possibilities are absurd or excluded, so $n = 3$ or 5.

If $n = 3$, each edge is incident with two flags fixed by r_1 , and \mathcal{M} is the second or third map in the second row of Figure 8 as these are transposed by r_0 or by r_0r_2 . If $n = 5$, each edge is incident with one flag fixed by r_1 , so $\text{Aut } \mathcal{M}$ acts regularly on the edges and thus sharply 2-homogeneously on the vertices. However, there is no sharply 2-homogeneous group of degree 5, so this case cannot arise. \square

ORCID iDs

Gareth A. Jones  <https://orcid.org/0000-0002-7082-7025>

References

- [1] N. Biggs, Classification of complete maps on orientable surfaces, *Rend. Mat. (6)* **4** (1971), 645–655.
- [2] R. P. Bryant and D. Singerman, Foundations of the theory of maps on surfaces with boundary, *Quart. J. Math. Oxford Ser. (2)* **36** (1985), 17–41, doi:10.1093/qmath/36.1.17.
- [3] H. S. M. Coxeter and W. O. J. Moser, *Generators and relations for discrete groups*, volume 14 of *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*, Springer-Verlag, Berlin-New York, 4th edition, 1980.
- [4] J. E. Graver and M. E. Watkins, Locally finite, planar, edge-transitive graphs, *Mem. Amer. Math. Soc.* **126** (1997), vi+75, doi:10.1090/memo/0601.
- [5] J. L. Gross and T. W. Tucker, *Topological graph theory*, Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, Inc., New York, 1987, a Wiley-Interscience Publication.

- [6] B. Grünbaum and G. C. Shephard, Edge-transitive planar graphs, *J. Graph Theory* **11** (1987), 141–155, doi:10.1002/jgt.3190110204.
- [7] L. Heffter, Ueber metacyklische Gruppen und Nachbarconfigurationen, *Math. Ann.* **50** (1898), 261–268, doi:10.1007/BF01448067.
- [8] B. Huppert, *Endliche Gruppen. I*, Die Grundlehren der Mathematischen Wissenschaften, Band 134, Springer-Verlag, Berlin-New York, 1967.
- [9] L. D. James, Imbeddings of the complete graph, *Ars Combin.* **16** (1983), 57–72.
- [10] L. D. James, Edge-symmetric orientable imbeddings of complete graphs, *European J. Combin.* **11** (1990), 133–144, doi:10.1016/S0195-6698(13)80067-4.
- [11] L. D. James and G. A. Jones, Regular orientable imbeddings of complete graphs, *J. Combin. Theory Ser. B* **39** (1985), 353–367, doi:10.1016/0095-8956(85)90060-7.
- [12] G. A. Jones, Automorphism groups of edge-transitive maps, 2019, arXiv:1605.09461v3 [math.CO], <https://arxiv.org/abs/1605.09461>.
- [13] G. A. Jones and J. S. Thornton, Operations on maps, and outer automorphisms, *J. Combin. Theory Ser. B* **35** (1983), 93–103, doi:10.1016/0095-8956(83)90065-5.
- [14] V. P. Korzhik and H.-J. Voss, On the number of nonisomorphic orientable regular embeddings of complete graphs, *J. Combin. Theory Ser. B* **81** (2001), 58–76, doi:10.1006/jctb.2000.1993.
- [15] S. Levy (ed.), *The eightfold way: the beauty of Klein's quartic curve*, volume 35 of *Mathematical Sciences Research Institute Publications*, Cambridge University Press, Cambridge, 1999.
- [16] C. H. Li, Vertex transitive embeddings of complete graphs, *J. Combin. Theory Ser. B* **99** (2009), 447–454, doi:10.1016/j.jctb.2008.09.002.
- [17] A. Orbanić, D. Pellicer, T. Pisanski and T. W. Tucker, Edge-transitive maps of low genus, *Ars Math. Contemp.* **4** (2011), 385–402, doi:10.26493/1855-3974.249.3a6.
- [18] D. Passman, *Permutation groups*, W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [19] D. Singerman, Klein's Riemann surface of genus 3 and regular imbeddings of finite projective planes, *Bull. London Math. Soc.* **18** (1986), 364–370, doi:10.1112/blms/18.4.364.
- [20] J. Širáň, T. W. Tucker and M. E. Watkins, Realizing finite edge-transitive orientable maps, *J. Graph Theory* **37** (2001), 1–34, doi:10.1002/jgt.1000.abs.
- [21] S. Wilson, Operators over regular maps, *Pacific J. Math.* **81** (1979), 559–568, <http://projecteuclid.org/euclid/pjm/1102785296>.
- [22] S. Wilson, Cantankerous maps and rotary embeddings of K_n , *J. Combin. Theory Ser. B* **47** (1989), 262–273, doi:10.1016/0095-8956(89)90028-2.
- [23] S. Wilson, Edge-transitive maps and non-orientable surfaces, *Math. Slovaca* **47** (1997), 65–83, proceedings of a meeting of GEMS, Graph embeddings and maps on surfaces (Donovaly, 1994).
- [24] H. Zassenhaus, Kennzeichnung endlicher linearer Gruppen als Permutationsgruppen, *Abh. Math. Sem. Univ. Hamburg* **11** (1935), 17–40, doi:10.1007/BF02940711.