

An infinite family of incidence geometries whose incidence graphs are locally X^*

Natalia Garcia-Colin

*CONACYT-INFOTEC. Circuito Tecnopolo Sur No 112, Col. Fracc. Tecnopolo Pocitos
C.P. 20313, Aguascalientes, Ags. México*

Dimitri Leemans[†] 

*Université Libre de Bruxelles, Département de Mathématique
C.P.216 - Algèbre et Combinatoire, Boulevard du Triomphe, 1050 Brussels, Belgium*

Received 16 September 2019, accepted 3 February 2020, published online 13 August 2021

Abstract

We construct a new infinite family of incidence geometries of arbitrarily large rank. These geometries are thick and residually connected and their type-preserving automorphism groups are symmetric groups. We also compute their Buekenhout diagram. The incidence graphs of these geometries are locally X graphs, but more interestingly, the automorphism groups act transitively, not only on the vertices, but more strongly on the maximal cliques of these graphs.

DEDICATED TO THE MEMORY OF BRANKO GRÜNBAUM.

Keywords: Kneser graph, locally X graph, incidence geometry.

Math. Subj. Class.: 05C75, 51E24

The influence of Branko in the world combinatorics is unbounded. Although the authors of this work did not have the luck of meeting him in person, they have being inspired by him through his influential writings on the theories of Convex Polytopes and Configurations of Points and Lines.

This paper contributes to the study of incidence geometries whose type-preserving automorphism groups are symmetric groups, this subject possesses a straight-forward connection to Branko's work on point-line configurations, the latter being incidence geometries too.

*The authors thank an anonymous referee for useful comments on the first version of this paper.

[†]Corresponding author.

E-mail addresses: natalia.garcia@infotec.mx (Natalia Garcia-Colin), Leemans.Dimitri@ulb.be (Dimitri Leemans)

1 Introduction

In [9, 7], Francis Buekenhout, Philippe Cara and Michel Dehon determined the so-called inductively minimal geometries, that are the thin residually connected regular incidence geometries of rank $n - 1$ with type-preserving automorphism group S_n . Recently, Maria Elisa Fernandes and Dimitri Leemans, extended this study to rank $n - 2$ thin geometries [19], and Fernandes and Leemans together with Mark Mixer classified the thin residually connected regular incidence geometries of rank $n - i$ ($i = 1, \dots, 4$) with a linear Buekenhout diagram and type preserving automorphism group S_n [18, 20], these geometries being abstract regular polytopes. Here we start from Kneser graphs and construct thick geometries whose underlying incidence graphs turn out to be locally X graphs. This provides yet another example of the natural connections between incidence geometries and other combinatorial structures.

Our construction is as follows (see Section 2 for terminology and notation).

Construction 1.1. For any positive integer $r \geq 2$ and a given Kneser graph $KG(n, k)$, define a rank r incidence system $\Gamma(KG(n, k), r) := (Y, *, t, I)$ as follows. Let $\Omega := \{1, \dots, n + k(r - 2)\}$. Let $I := \{1, \dots, r\}$. Take r copies Y_1, \dots, Y_r of the set of all the subsets of size k of Ω , and let $Y = Y_1 \cup \dots \cup Y_r$. For any $x \in Y_i$, define $t(x) = i$. For any elements $x_i \in Y_i$ and $x_j \in Y_j$, we say that $x_i * x_j$ if and only if $i \neq j$ and x_i and x_j are disjoint as subsets of Ω .

These geometries have several nice properties summarized as follows.

Theorem 1.2. $\Gamma(KG(n, k), r)$ is a thick, residually connected and flag-transitive incidence geometry of rank r . Its automorphism groups are $\text{Aut}_I(\Gamma) \cong S_{n+k(r-2)}$ and $\text{Aut}(\Gamma) \cong S_{n+k(r-2)} \times S_r$. The Buekenhout diagram of $\Gamma(KG(n, k), r)$ is a complete graph. The orders of the diagram are $\binom{n-k}{k-1}$, the number of elements of each type is $\binom{n+k(r-2)}{k}$, the edges are labelled as $d - g - d$ where $g = 3$ if $n = 2k + 1$ and $g = 2$ otherwise, and $d = 2\lceil \frac{k}{n-2k} \rceil + 1$.

We want to highlight one remarkable member of this infinite family. The geometry $\Gamma(KG(5, 2), r)$ has arbitrary large rank r and all of its rank two residues are isomorphic to the Desargues configuration.

We found these new geometries while searching for large locally X graphs. For a given graph X , a graph \mathcal{G} is locally X if the graphs induced on the neighbours of every vertex of \mathcal{G} are isomorphic to X . In the literature, \mathcal{G} is also referred to as an *extension of X* or a *locally homogeneous graph*.

The construction of the incidence graph corresponding to the incidence geometry in Theorem 1.2 immediately gives the following corollary, adding explicitly one large family to the list of known locally X graphs.

Corollary 1.3. The incidence graph of $\Gamma(KG(n, k), r)$ is a locally $KG(n + k(r - 3), k) \times K_{r-1}$ graph.

These graphs are isomorphic to $KG(n + k(r - 2), k) \times K_r$, where \times stands for the tensor product. Thus their vertex-transitivity is easy to prove, implying that they are locally $KG(n + k(r - 3), k) \times K_{r-1}$. However, Theorem 1.2 proves more than the above, namely:

Corollary 1.4. The symmetric group $S_{n+k(r-2)}$ acts transitively on the set of maximal cliques of $KG(n + k(r - 2), k) \times K_r$.

We believe that the connection between incidence geometries and locally X graphs is worthy of further study. It is interesting to explore under which conditions incidence geometries can lead to locally X graphs.

The paper is organized as follows. In Section 2 we present some structural properties of Kneser graphs which will be used in the construction of the incidence geometries we study and we present the necessary background on incidence geometries. In Section 3 we compute the neighbourhood geometry of a Kneser graph, which is used in Section 4 to construct a new infinite family of locally X graphs.

2 Preliminaries

2.1 Graph theory

A Kneser graph¹ $KG(n, k)$ is a graph whose vertex set is the set of all k -subsets of $\{1, \dots, n\}$ and any pair of disjoint k -subsets is joined by an edge. Some Kneser graphs are very familiar objects, for instance the complete graphs, K_n , are Kneser graphs $KG(n, 1)$ and the Petersen graph is a $KG(5, 2)$. Its very simple to see that Kneser graphs are locally Kneser graphs, furthermore Jonathan Hall proved in [21] that there are exactly three pairwise non-isomorphic locally Petersen graphs, only one of them being a Kneser graph, namely $KG(7, 2)$.

The following lemmas will cover some structural characteristics of Kneser graphs which, in turn, will be used in the later sections for determining structural characteristics of our constructions.

Lemma 2.1. *The smallest odd cycle of a Kneser graph, $KG(n, k)$, with $n > 2k + 1$ has length $2\lceil \frac{k}{n-2k} \rceil + 1$.*

Proof. We may assume that $n < 3k$, as $KG(n, k)$ with $n \geq 3k$ has triangles, and the statement holds. Let $n = 2k + r$ for some $r < k$ and A_1, \dots, A_{2l+1} be the vertices of the smallest odd cycle, as subsets of $[n]$.

By construction, we have $A_1 \cap A_2 = \emptyset$ and $A_2 \cap A_3 = \emptyset$ thus $A_1 \cup A_3 \subset A_2^c$ and they have non empty intersection as, $|A_2^c| = n - k = k + r < 2k$, thus $|A_1 \cap A_3| \geq k - r$ and A_1 and A_3 cannot be adjacent. Similarly, we can argue that $|A_3 \cap A_5| \geq k - r$ thus $|A_1 \cap A_5| \geq k - 2r$. We may continue this process to conclude that $|A_1 \cap A_{2i+1}| \geq k - ir$.

Hence $A_1 \cap A_{2l+1} = \emptyset$ if and only if $k - lr \leq 0$. This happens precisely when $\lceil \frac{k}{n-2k} \rceil \leq l$ and the result follows. \square

Lemma 2.2. *Between any two vertices of a Kneser graph, A, B , such that $|A \cap B| = c$ there is an even path of length $2\lceil \frac{k-c}{n-2k} \rceil$ and an odd path of length $2\lceil \frac{c}{n-2k} \rceil + 1$.*

Proof. Let A and B be two vertices of $KG(n, k)$, $C = A \cap B$, $|C| = c$, $D = (A \cup B)^c$, and $|D| = n - 2k + c$. Let X_1, \dots, X_l and Y_1, \dots, Y_l with $l = \lceil \frac{k-s}{n-2k} \rceil$ be partitions of $A \setminus B$ and $B \setminus A$, respectively, in sets of size $n - 2k$, perhaps except the last one.

Let $A_{2i-1} = (\cup_{j=1}^i X_j) \cup (\cup_{j=i+1}^l Y_j) \cup D'$ for some $D' \subset D$ of cardinality c and $A_{2i} = (\cup_{j=1}^i Y_j) \cup (\cup_{j=i+1}^l X_j) \cup C$ for $0 \leq i \leq l$. Clearly $A_j \cap A_{j+1} = \emptyset$, $A = A_0$, $B = A_{2l}$, thus $A, A_1, \dots, A_{2l-1}, B$ is a path of even length $2\lceil \frac{k-c}{n-2k} \rceil$.

For the odd path, take $D' \subset D$ of cardinality c and let $A' = B \setminus A \cup D'$. Then A and A' are adjacent, and we may construct an even path of size $2\lceil \frac{c}{n-2k} \rceil$ between A' and B as before, given that $|A' \cap B| = k - c$, and the result follows. \square

¹Lovász introduced the term Kneser graph in [30] after a problem posed by Kneser in [27].

2.2 Incidence geometry

An *incidence system* [8], $\Gamma := (Y, *, t, I)$ is a 4-tuple such that

- Y is a set whose elements are called the *elements* of Γ ;
- I is a set whose elements are called the *types* of Γ ;
- $t : Y \rightarrow I$ is a *type function*, associating to each element $x \in Y$ of Γ a type $t(x) \in I$;
- $*$ is a binary relation on Y called *incidence*, that is reflexive, symmetric and such that for all $x, y \in Y$, if $x * y$ and $t(x) = t(y)$ then $x = y$.

The *incidence graph* of Γ is the graph whose vertex set is Y and where two vertices are joined provided the corresponding elements of Γ are incident.

A *flag* is a set of pairwise incident elements of Γ , i.e. a clique of its incidence graph. The *type* of a flag F is $\{t(x) : x \in F\}$. A *chamber* is a flag of type I . An element x is *incident* to a flag F and we write $x * F$ for that, when x is incident to all elements of F . An incidence system Γ is a *geometry* or *incidence geometry* if every flag of Γ is contained in a chamber (or in other words, every maximal clique of the incidence graph is a chamber). The *rank* of Γ is the number of types of Γ , namely the cardinality of I .

Observe that the incidence graph of an incidence system of rank n is an n -partite graph. This will play a key role in the construction of an infinite family of locally X graphs.

Let $\Gamma := (Y, *, t, I)$ be an incidence system. Given $J \subseteq I$, the J -*truncation* of Γ is the incidence system $\Gamma^J := (t^{-1}(J), *|_{t^{-1}(J) \times t^{-1}(J)}, t|_J, J)$. In other words, it is the subgeometry constructed from Γ by taking only elements of type J and restricting the type function and incidence relation to these elements.

Let $\Gamma := (Y, *, t, I)$ be an incidence system. Given a flag F of Γ , the *residue* of F in Γ is the incidence system $\Gamma_F := (Y_F, *_F, t_F, I_F)$ where

- $Y_F := \{x \in Y : x * F, x \notin F\}$;
- $I_F := I \setminus t(F)$;
- t_F and $*_F$ are the restrictions of t and $*$ to Y_F and I_F .

An incidence system Γ is *residually connected* when each residue of rank at least two of Γ has a connected incidence graph. It is called *firm* (resp. *thick*) when every residue of rank one of Γ contains at least two (resp. three) elements.

Let $\Gamma := (Y, *, t, I)$ be an incidence system. An *automorphism* of Γ is a mapping $\alpha : (Y, I) \rightarrow (Y, I) : (x, t(x)) \mapsto (\alpha(x), t(\alpha(x)))$ where


- α is a bijection on Y ;
- for each $x, y \in Y$, $x * y$ if and only if $\alpha(x) * \alpha(y)$;
- α induces a bijection on I such that for each $x, y \in Y$, $t(x) = t(y)$ if and only if $t(\alpha(x)) = t(\alpha(y))$.

An automorphism α of Γ is called *type preserving* when for each $x \in Y$, $t(\alpha(x)) = t(x)$. The set of all automorphisms of Γ together with the composition forms a group that is called the *automorphism group* of Γ and denoted by $Aut(\Gamma)$. The set of all type-preserving automorphisms of Γ is a subgroup of $Aut(\Gamma)$ that we denote by $Aut_I(\Gamma)$. An incidence system Γ is *flag-transitive* if $Aut_I(\Gamma)$ is transitive on all flags of a given type J for each type $J \subseteq I$.

A rank two geometry with points and lines is called a *generalised digon* if every point is incident to every line and, it is called a *partial linear space* if there is at most one line through any pair of points. An incidence geometry is said to satisfy the *intersection property of rank 2*, denoted by $(IP)_2$, when all its rank two residues are either partial linear spaces or generalised digons.

Let Γ be a firm, residually connected and flag-transitive geometry. The *Buekenhout diagram* of Γ is a graph whose vertices are the elements of I and with an edge $\{i, j\}$ with label $d_{ij} - g_{ij} - d_{ji}$ whenever every residue of type $\{i, j\}$ is not a generalised digon. The number g_{ij} is called the *gonality* and is equal to half the girth of the incidence graph of a residue of type $\{i, j\}$. The number d_{ij} is called the *i-diameter* of a residue of type $\{i, j\}$ and is the longest distance from an element of type i to any element in the incidence graph of a residue of type $\{i, j\}$. Moreover, to every vertex i is associated a number s_i , called the *i-order*, which is equal to the size of a residue of type i minus one, and a number n_i which is the number of elements of type i of the geometry.


The Petersen graph, for instance, can be seen as a geometry of rank two whose elements are the vertices and edges of the graph. Its Buekenhout diagram is the following.



Let G be a graph. Denote its set of vertices (resp. edges) by G_0 (resp. G_1). For distinct $p, q \in G_0$, we say that p and q are *adjacent* – and we write $p \sim q$ – whenever $\{p, q\} \in G_1$. As in [29], to the graph G , we associate a new rank 2 geometry \tilde{G} , called the *neighborhood geometry* of G , whose elements are, roughly speaking, the vertices and the neighborhoods of vertices of G . More precisely, we define \tilde{G} to be the geometry $(G_0 \times \{0\} \cup G_0 \times \{1\}, \tilde{*}, \tilde{t}, \{0, 1\})$ with

- $\tilde{t}(G_0 \times \{i\}) = i$, for $i = 0, 1$;
- $(p, 0)\tilde{*}(q, 1)$ iff $p \sim q$, for $p, q \in G_0$.

As pointed out in [29, Table 1], the neighborhood geometry of the Petersen graph is Desargues’ configuration. Figure 1 gives the Petersen graph and Desargues’ configuration. The Buekenhout diagram of Desargues’ configuration is the following.



3 The neighborhood geometry of a Kneser graph

In this section, we compute the neighborhood geometry of a given Kneser graph. These geometries will then be used in the next section to construct locally X graphs as incidence graphs of some particular incidence geometries. The incidence graphs of these geometries are sometimes called the bipartite Kneser graphs.




Figure 1: Petersen graph $KG(5, 2)$ and Desargues' configuration $K\tilde{G}(5, 2)$

Lemma 3.1. *The 0-diameter and 1-diameter of $K\tilde{G}(n, k)$ is $2\lceil \frac{k}{n-2k} \rceil + 1$.*

Proof. As the construction of $K\tilde{G}(n, k)$ is symmetric in the set of types, the 0-diameter and the 1-diameter are the same.

By Lemma 2.2 the distance between any two vertices of $K\tilde{G}(n, k)$ is at most $2\lceil \frac{k-c}{n-2k} \rceil$ or $2\lceil \frac{c}{n-2k} \rceil + 1$, for some $0 \leq c \leq k$. This achieves a maximum of $2\lceil \frac{k}{n-2k} \rceil + 1$ when $c = k$, that is, when we are tracing a path from the two copies of the same vertex in $K\tilde{G}(n, k)$.

We now argue that this bound can't be improved, as such improvement would contradict Lemma 2.1. □

Lemma 3.2. *The gonality of $K\tilde{G}(n, k)$ is 3 when $n = 2k + 1$ and 2 when $n \geq 2k + 2$.*

Proof. If $n = 2k + 1$, the following is a circuit of length 6 in the incidence graph of $K\tilde{G}(n, k)$: $(\{1, \dots, k\}, 0) \tilde{*} (\{k+1, \dots, 2k\}, 1) \tilde{*} (\{1, \dots, k-1, 2k+1\}, 0) \tilde{*} (\{k, \dots, 2k-1\}, 1) \tilde{*} (\{1, \dots, k-1, 2k\}, 0) \tilde{*} (\{k+1, \dots, 2k-1, 2k+1\}, 1) \tilde{*} (\{1, \dots, k\}, 0)$.

If $n > 2k + 1$, the following is a circuit of length 4 in the incidence graph of $K\tilde{G}(n, k)$: $(\{1, \dots, k\}, 0) \tilde{*} (\{k+1, \dots, 2k\}, 1) \tilde{*} (\{1, \dots, k-1, 2k+1\}, 0) \tilde{*} (\{k+1, \dots, 2k-1, 2k+2\}, 1) \tilde{*} (\{1, \dots, k\}, 0)$. □

Lemma 3.3. *$K\tilde{G}(n, k)$ is a connected graph.*

Proof. This follows from the fact that $KG(n, k)$ has cycles of odd length as proven in Lemma 2.1. □

4 Proof of Theorem 1.2

In this section we prove the main theorem of this paper. Observe that the case $r = 2$ in Construction 1.1 gives the neighbourhood geometry $K\tilde{G}(n, k)$ of the Kneser graph $KG(n, k)$ that was defined in the previous section.

Lemma 4.1. *$\Gamma(KG(n, k), r)$ is an incidence geometry.*

Proof. As $|\Omega| = n + k(r - 2)$ and $n \geq 2k + 1$, it is always possible to find r pairwise disjoint subsets of size k in Ω . Hence every maximal flag of Γ must be a chamber and Γ is an incidence geometry. □

Lemma 4.2. *The symmetric group $S_\Omega \cong S_{n+k(r-2)}$ acts transitively on the chambers of $\Gamma(KG(n, k), r)$ (or in other words, $\Gamma(KG(n, k), r)$ is flag-transitive).*

Proof. This is due to the fact that S_Ω is $(n + k(r - 2))$ -transitive on Ω . □

Lemma 4.3. *$\Gamma(KG(n, k), r)$ is residually connected.*

Proof. We prove this by induction on r . The case $r = 2$ is dealt with in Lemma 3.3.

Suppose that $\Gamma(KG(n, k), r)$ is residually connected. In order to prove that $\Gamma(KG(n, k), r + 1)$ is residually connected, we only need to show that the incidence graph of $\Gamma(KG(n, k), r + 1)$ is connected, as all residues of $\Gamma(KG(n, k), r + 1)$ of rank $< r + 1$ are connected by the induction hypothesis and the fact that $\Gamma(KG(n, k), r + 1)$ is flag-transitive as shown in Lemma 4.2. Take x_i an element of type i and x_j an element of type $j \neq i$. The $\{i, j\}$ -truncation of $\Gamma(KG(n, k), r)$ is the neighborhood geometry $\tilde{K}G(n * k(r - 2), k)$ of a Kneser graph $KG(n * k(r - 2), k)$. By Lemma 3.3, $\tilde{K}G(n * k(r - 2), k)$ is connected. Hence, every rank two truncation of $\Gamma(KG(n, k), r)$ is connected and therefore $\Gamma(KG(n, k), r)$ is connected. □

Lemma 4.4. *For any $i = 1, \dots, r$, the i -order of $\Gamma(KG(n, k), r)$ is equal to $\binom{n-k}{k-1}$.*

Proof. A flag F of rank $r - 1$ consists of $r - 1$ pairwise disjoint subsets of size k . Hence these subsets cover $k(r - 1)$ points of Ω . So there are $n + k(r - 2) - k(r - 1) = n - k$ points not covered by F in Ω . There are thus $\binom{n-k}{k}$ subsets of size k that are disjoint with all subsets of F . □

The previous lemma immediately implies the following corollary.

Corollary 4.5. *$\Gamma(KG(n, k), r)$ is thick.*

Lemma 4.6. *Every rank two residue of $\Gamma(KG(n, k), r)$ is isomorphic to $\tilde{K}G(n, k)$.*

Proof. Every rank two residue is obtained from a flag F that has $r - 2$ elements. These $r - 2$ elements cover $k(r - 2)$ elements of Ω and therefore there are n elements of Ω remaining. □

Lemma 4.7. *$\text{Aut}_I(\Gamma(KG(n, k), r)) \cong S_{n+k(r-2)}$ and $\text{Aut}(\Gamma(KG(n, k), r)) \cong S_{n+k(r-2)} \times S_r$*

Proof. By Lemma 4.2, we know that $\text{Aut}_I(\Gamma(KG(n, k), r)) \geq S_{n+k(r-2)}$. Obviously, it cannot be strictly bigger. Construction 1.1 is symmetric in the set of types. Hence $\text{Aut}(\Gamma(KG(n, k), r)) \cong S_{n+k(r-2)} \times S_r$. □

Theorem 1.2 is a summary of all the properties we have proved on $\Gamma(KG(n, k), r)$ above. Note that the gonality and diameters of the rank two residues of the Buekenhout diagram were computed in Lemma 3.2 and Lemma 3.1.

We highlight that $\Gamma(KG(5, 2), 3)$ was already given in [10, page 86]. Furthermore, geometries satisfying the intersection property of rank two, $(IP)_2$, attracted much attention in the nineties and noughties² (see, for instance, [10, 28]). It turns out that some of the incidence geometries obtained by Construction 1.1 satisfy $(IP)_2$, as shown in the next corollary.

²The noughties mean the years 2000–2010.

Corollary 4.8. $\Gamma(KG(n, k), r)$ is $(IP)_2$ if and only if $n = 2k + 1$.

Proof. For a rank two residue of $\Gamma(KG(n, k), r)$ to satisfy $(IP)_2$, we need it to be a generalised digon or its gonality to be at least 3. Generalised digons have gonality and diameters equal to two. Lemma 3.1 and Lemma 3.2 then finish the proof. \square

Observe that when $n = 2k + 1$, the diameters and gonality written on the edges of the Buekenhout diagram are respectively n and 3.

5 Connection with locally X graphs

A. Zykov [38, 39] posed the problem of characterizing the graphs, X , for which there are locally X graphs. Finding any general solution for this problem is difficult, if at all possible.

Apart from the inherent interest of this problem for graph theorists, another motivation for the study of locally homogeneous graphs is observed in [23]:

“The theorems may find application in the characterization of the Johnson scheme among the primitive association schemes and distance regular graphs. It can also be used to characterize alternating and symmetric groups (of sufficiently large degree) by centralizers of various of their elements (the initial motivation for the theorem).”

The progress thus far has followed three general lines of enquiry; the undecidability of the problem; the construction of locally X graphs for some selected graphs, X ; and sufficient conditions for a graph X to have an extension. Our construction adds to the second line. We now prove Corollary 1.3.

Proof of Corollary 1.3. Since $\Gamma(KG(n, k), r)$ is a flag-transitive geometry and since $\text{Aut}(\Gamma(KG(n, k), r)) \cong S_{n+k(r-2)} \times S_r$, all the incidence graphs of the residues of rank $r - 1$ are isomorphic. \square

Corollary 5.1. *There exists a locally Desargues graph whose automorphism group is isomorphic to S_7 .*

Proof. Such a graph can be obtained as the incidence graph of $\Gamma(KG(5, 2), 3)$. \square

It would be interesting to explore what are the characteristics of incidence geometries that lead to interesting locally X graphs.

We would like to highlight that in [22], a complete list of all graphs X of order up to six having an extension is given; in some cases all such extensions are characterized. Constructions of locally X graphs, for instance, cycles [16, 15], unions of paths [34], trees [2], polyhedra [3, 13, 14], the Petersen graph [21], other Kneser graphs [23, 32], dense graphs [12, 33] and others have been investigated in [4, 5, 6, 11, 17, 24, 26, 37]. A rich compilation of locally X graphs can also be found in [36]. Some structural characteristics for a graph X to have an extension have been given in [1, 5, 6, 25, 35].

Finally, as pointed out by a referee, there are also lots of interesting examples of locally X graphs that come from abstract regular polytopes. For example, as mentioned in [31, page 165], if \mathcal{P} is a regular n -polytope with triangular faces, the 1-skeleton (or edge graph) of \mathcal{P} is a locally X graph where X is the 1-skeleton of the vertex-figure of \mathcal{P} . Every polytope with a Schläfli symbol of the form $\{3, p_2, \dots, p_n\}$ is of this kind. For example, there are lots of interesting locally X graph where X is the graph of a toroidal map (and these are obtained from polytopes of type $\{3, 4, 4\}$, $\{3, 3, 6\}$ or $\{3, 6, 3\}$).

ORCID iDs

Dimitri Leemans  <https://orcid.org/0000-0002-4439-502X>

References

- [1] S. Al-Addasi, Some properties of locally homogeneous graphs, *An. Ştiinţ. Univ. "Ovidius" Constanţa Ser. Mat.* **18** (2010), 15–21, <https://www.anstuocmath.ro/volume-xviii-2010-fascicola-2.html>.
- [2] A. Blass, F. Harary and Z. Miller, Which trees are link graphs?, *J. Combin. Theory Ser. B* **29** (1980), 277–292, doi:10.1016/0095-8956(80)90085-4.
- [3] A. Blokhuis, A. E. Brouwer, D. Buset and A. M. Cohen, The locally icosahedral graphs, in: *Finite geometries (Winnipeg, Man., 1984)*, Dekker, New York, volume 103 of *Lecture Notes in Pure and Appl. Math.*, pp. 19–22, 1985.
- [4] A. E. Brouwer, J. H. Koolen and M. H. Klin, A root graph that is locally the line graph of the Petersen graph, volume 264, pp. 13–24, 2003, doi:10.1016/s0012-365x(02)00546-0, the 2000 *Com²MaC* Conference on Association Schemes, Codes and Designs (Pohang).
- [5] M. Brown and R. Connelly, On graphs with a constant link, in: *New directions in the theory of graphs (Proc. Third Ann Arbor Conf., Univ. Michigan, Ann Arbor, Mich., 1971)*, 1973 pp. 19–51.
- [6] M. Brown and R. Connelly, On graphs with a constant link. II, *Discrete Math.* **11** (1975), 199–232, doi:10.1016/0012-365x(75)90037-0.
- [7] F. Buekenhout and P. Cara, Some properties of inductively minimal geometries, *Bull. Belg. Math. Soc. - Simon Stevin* **5** (1998), doi:10.36045/bbms/1103409005.
- [8] F. Buekenhout and A. M. Cohen, *Diagram Geometry. Related to Classical Groups and Buildings.*, volume 57 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, Springer-Verlag Berlin Heidelberg, 2013, doi: 10.1007/978-3-642-34453-4.
- [9] F. Buekenhout, M. Dehon and P. Cara, Inductively minimal flag-transitive geometries, in: *Mostly finite geometries (Iowa City, IA, 1996)*, Dekker, New York, volume 190 of *Lecture Notes in Pure and Appl. Math.*, pp. 185–190, 1997.
- [10] F. Buekenhout, M. Dehon and D. Leemans, An atlas of residually weakly primitive geometries for small groups, *Acad. Roy. Belg. Cl. Sci. Mém. Collect. 8° (3)* **14** (1999), 175.
- [11] F. Buekenhout and X. Hubaut, Locally polar spaces and related rank 3 groups, *J. Algebra* **45** (1977), 391–434, doi:10.1016/0021-8693(77)90334-9.
- [12] P. Bugata, M. Horňák and S. Jendroľ, On graphs with given neighbourhoods, *Časopis Pěst. Mat.* **114** (1989), 146–154, doi:10.21136/cpm.1989.108714.
- [13] D. Buset, Graphs which are locally a cube, *Discrete Math.* **46** (1983), 221–226, doi:10.1016/0012-365x(83)90116-4.
- [14] D. Buset, Locally polyhedral graphs, in: *Finite geometries (Winnipeg, Man., 1984)*, Dekker, New York, volume 103 of *Lecture Notes in Pure and Appl. Math.*, pp. 23–25, 1985.
- [15] D. Buset, Locally C_n^k graphs, *Bull. Belg. Math. Soc. Simon Stevin* **2** (1995), 481–485, <http://projecteuclid.org/euclid.bbms/1103408672>.
- [16] B. L. Chilton, R. Gould and A. D. Polimeni, A note on graphs whose neighborhoods are n -cycles, *Geometriae Dedicata* **3** (1974), 289–294, doi:10.1007/bf00181321.

- [17] J. Doyen, X. Hubaut and M. Reynaert, Finite graphs with isomorphic neighbourhood, *Problèmes Combinatoires et Théorie des Graphes, Colloque C.N.R.S., Orsay (1976)*.
- [18] M. E. Fernandes and D. Leemans, Polytopes of high rank for the symmetric groups, *Adv. Math.* **228** (2011), 3207–3222, doi:10.1016/j.aim.2011.08.006.
- [19] M. E. Fernandes and D. Leemans, C-groups of high rank for the symmetric groups, *J. Algebra* **508** (2018), 196–218, doi:10.1016/j.jalgebra.2018.04.031.
- [20] M. E. Fernandes, D. Leemans and M. Mixer, An extension of the classification of high rank regular polytopes, *Trans. Amer. Math. Soc.* **370** (2018), 8833–8857, doi:10.1090/tran/7425.
- [21] J. I. Hall, Locally Petersen graphs, *J. Graph Theory* **4** (1980), 173–187, doi:10.1002/jgt.3190040206.
- [22] J. I. Hall, Graphs with constant link and small degree or order, *J. Graph Theory* **9** (1985), 419–444, doi:10.1002/jgt.3190090313.
- [23] J. I. Hall, A local characterization of the Johnson scheme, *Combinatorica* **7** (1987), 77–85, doi:10.1007/bf02579203.
- [24] J. I. Hall and E. E. Shult, Locally cotriangular graphs, *Geom. Dedicata* **18** (1985), 113–159, doi:10.1007/bf00151394.
- [25] P. Hell, Graphs with given neighborhoods. I, in: *Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976)*, CNRS, Paris, volume 260 of *Colloq. Internat. CNRS*, pp. 219–223, 1978.
- [26] P. Hell, H. Levinson and M. Watkins, Some remarks on transitive realizations of graphs., in: *Proc. 2nd. Car. Conf. Comb. and Comp.*, 1977 pp. 115–122.
- [27] M. Kneser, Aufgabe 360, *Jahresber. DMV* **58** (1956).
- [28] D. Leemans, *Residually weakly primitive and locally two-transitive geometries for sporadic groups*, Publication de la Classe des sciences. Collection in-4 3. sér., t. 11., Classe des sciences, Académie royale de Belgique, 2008, 2008.
- [29] C. Lefèvre-Percsy, N. Percsy and D. Leemans, New geometries for finite groups and polytopes, *Bull. Belg. Math. Soc. Simon Stevin* **7** (2000), 583–610, <http://projecteuclid.org/euclid.bbms/1103055618>.
- [30] L. Lovász, Kneser’s conjecture, chromatic number, and homotopy, *J. Combin. Theory Ser. A* **25** (1978), 319–324, doi:10.1016/0097-3165(78)90022-5.
- [31] P. McMullen and E. Schulte, *Abstract regular polytopes*, volume 92 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, Cambridge, 2002, doi:10.1017/cbo9780511546686.
- [32] A. Moon, The graphs $G(n, k)$ of the Johnson schemes are unique for $n \geq 20$, *J. Combin. Theory Ser. B* **37** (1984), 173–188, doi:10.1016/0095-8956(84)90070-4.
- [33] R. Nedela, Locally homogeneous graphs with dense links at vertices, *Czechoslovak Math. J.* **42(117)** (1992), 515–517.
- [34] T. D. Parsons and T. Pisanski, Graphs which are locally paths, in: *Combinatorics and graph theory (Warsaw, 1987)*, PWN, Warsaw, volume 25 of *Banach Center Publ.*, pp. 127–135, 1989.
- [35] G. M. Weetman, A construction of locally homogeneous graphs, *J. London Math. Soc. (2)* **50** (1994), 68–86, doi:10.1112/jlms/50.1.68.
- [36] E. W. Weisstein, Local graph, From MathWorld—A Wolfram Web Resource, 2018, <https://mathworld.wolfram.com/LocalGraph.html>.
- [37] B. Zelinka, The least connected non-vertex-transitive graph with constant neighbourhoods, *Czechoslovak Math. J.* **40(115)** (1990), 619–624.

- [38] A. Zykov, Problem 30, in: M. Fielder (ed.), *Theory of graphs and its applications*, 1963 pp. 164–165.
- [39] A. A. Zykov, Graph-theoretical results of Novsibirsk mathematicians, in: *Theory of Graphs and its Applications (Proc. Sympos. Smolenice, 1963)*, Publ. House Czechoslovak Acad. Sci., Prague, 1963 pp. 151–153.